

**POISON**

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**TRICLOPYR**

**600EC** HERBICIDE

**ACTIVE CONSTITUENT:**

600 g/L TRICLOPYR  
present as the butoxyethyl ester

**GROUP I HERBICIDE**

For the control of a range of woody weeds and  
melons as specified in the Directions for Use.

**IMPORTANT: READ THIS BOOKLET BEFORE USE**  
**SHAKE WELL BEFORE USE**

**RELYON**

APVMA Approval No:  
89668/125736

## DIRECTIONS FOR USE

### RESTRAINTS:

- **DO NOT** apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result.
- **DO NOT** spray if rain is likely within one hour or if foliage is wet from rain and dew. However, when tankmixed with Panzer 450 this time extends to 6 hours.
- **DO NOT** burn off, cut or clear blackberry or other woody weeds for six months after spraying.

### 1. WOODY WEED SITUATIONS

**Table A: High Volume Spraying**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>			
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /100 L water</b>
African boxthorn	Less than 2m tall	Tas only	170 mL
Angophora and Banksia regrowth	1 to 2 m tall	All States	400 or 560 mL
Blackberry In association with: St John's wort	Active growth during late spring to early autumn		
	During flowering (Nov-Jan)	NSW, Vic and Tas only	
Brigalow	1 to 2 m tall	NSW and Qld only	
Broom: English	Spring to mid- summer prior to pod formation	All States	
Camphor laurel	Seedlings up to 3 m tall		
Capeweed	Rosette	Tas only	80 mL
Common prickly pear	Active growth	All States	3 L
English ivy	Active growth during late spring to late summer	Vic only	1 L + 1 L Glyphosate (360g/ L)
Eucalyptus	Seedlings and regrowth from small lignotubers, 1 to 2 m tall	Qld, SA, WA, and NT only	400 mL
		NSW, Tas and Vic only	560 mL
Fennel	1 to 2 m tall	Tas only	170 mL
Green cestrum		NSW, Qld and Vic only	

<b>CRITICAL COMMENTS</b>
Use the higher rate on larger regrowth. Ensure the weed has dense foliage.
Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before spraying with RELYON TRICLOPYR 600EC HERBICIDE. Any subsequent regrowth and seedlings should be sprayed after hardening off.
Apply as a thorough foliage spray.
Use at least 1000 L of water /ha.
<b>DO NOT</b> treat ivy growing up trees or on other plants as death of the host may result. This mixture is not selective to grasses.
Add Decol® S35B Surfactant at 250 mL to 500 mL/100 L of water for best results.
Some regrowth may be expected the following season which can be sprayed after hardening off.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /100 L water
Groundsel bush ( <i>Baccharis halimifolia</i> )	Seedlings, 1 to 2 m tall	All States	160 mL
	2 to 3 m tall		320 mL
Gorse	Spring to mid summer		170 mL or 340 mL
Horehound	Rosette	Tas only	170 mL
Saffron thistle	Up to bud stage		80 mL
Tiger pear	Active growth	All States	3 L
Wattles, including Silver wattle	Seedlings, 1 to 2 m tall		160 mL
Black wattle	2 to 3 m tall		320 mL

**Table B: Aerial Application**

**Helicopter NSW, SA, Tas, Vic and WA only**

**Helicopter or fixed wing aircraft (Qld only)**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha
Blackberry	Late spring to autumn	All States	4.8 L

**Table C: Controlled Droplet Application (C.D.A.)**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /1L water
Blackberry	Late spring to autumn	All States	170 mL

CRITICAL COMMENTS
Add a 100% concentrate non- ionic wetting agent at rate of 125 mL/100L water. Retreatment of regrowth may be necessary. Use higher water rate on older hardened off plants.

CRITICAL COMMENTS
AVOID overspray/drift onto waterways.

CRITICAL COMMENTS

**Table D: Low Volume High Concentrate Application Techniques  
(Gas Powered Gun, Sprinkler Sprayer)**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>			
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /10 L water</b>
Blackberry	Late spring to autumn	All States	280 mL
Eucalypt seedlings	1 to 2 m tall		400 mL

<b>CRITICAL COMMENTS</b>

**Table E: Basal Bark and Cut Stump Treatment**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>			
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /60 L diesel</b>
African boxthorn	<b>Basal bark:</b> Plants up to 5 cm basal diameter	All States	2 L
Australian blackthorn			1 L
Broom	<b>Cut stump:</b> Plants up to and in excess of basal bark sizes	Tas only	1.25 L
Brown salwood		All States	0.5 L
Bitter bark		NSW and Qld only	1 L
Castor oil Chinee apple		All States	
Dawson gum		Qld only	2 L
Eucalypts (except Dawson gum) False sandalwood Green wattle Lantana Needlewood	<b>Basal bark:</b> Plants up to 5 cm basal diameter  <b>Cut stump:</b> Plants up to and in excess of basal bark sizes	All States	1 L
Olive		SA only	2 L
Paperbark teatree Rubbervine Silver wattle		NSW only	4 L
Sweet briar		All States	1 L
Yellow-wood			2 L
		Qld only	

<b>CRITICAL COMMENTS</b>

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /60 L diesel
Camphor laurel	<b>Basal bark:</b> Plants up to 10 cm basal diameter  <b>Cut stump</b> Plants up to and in excess of basal bark sizes	NSW and Qld only	1 L
Common prickly pear		All States	0.8 L
Groundsel bush ( <i>Baccharis halimifolia</i> )			0.5 L
Prickly acacia			
Privet			5 L
Smooth tree pear			0.8 L
Tiger pear			
Tree of heaven			1 L
FENCELINES AND FIRE TRAILS only.			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /60 L diesel
Broadleaf hopbush, Narrowleaf hopbush, Turpentine bush	<b>Basal bark:</b> Plants up to 10 cm basal diameter	NSW only	1 L

## 2. CROPPING/FALLOW SITUATIONS

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

FALLOW, STUBBLE, FIREBREAKS.			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha
Camel (Afghan, Bitter) melon ( <i>Citrullus lanatus</i> )	Up to 20 cm diameter	NSW, Qld, SA, Vic and WA only	120 mL*
	Runners from 20 to 40 cm diameter		160 mL*
Prickly paddy melon ( <i>Cucumis myriocarpus</i> )	Up to 20 cm diameter		80 mL*
	Runners from 20 to 40 cm diameter		160 mL*

CRITICAL COMMENTS
Apply as a thorough foliage spray.
Treat from early summer rains to end of April when regrowth is apparent.
Treatment may be carried out at any time of the year.
Apply as a thorough foliage spray.
CRITICAL COMMENTS

CRITICAL COMMENTS
There are some strains of melon that are not controlled. Contact your Nutrien Ag Solutions representative for more information.
*Add a crop oil such as Uptake® Spraying Oil at 500 mL/100 L water or D-C-Trate® Crop Oil at 1 L/100 L water. <b>DO NOT</b> use oils when tank mixing with Panzer 450 Herbicide.
See <b>COMPATIBILITY</b> section. This mixture is not selective to grasses.
When using RELYON TRICLOPYR 600EC HERBICIDE and Panzer 450 by aerial application, observance of a buffer zone of 150 metres to protect native tree species is required.

SORGHUM (Apply between 4 to 6 leaf stage, when secondary roots have developed)			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha
Prickly paddy melon	Up to 20 cm diameter	NSW and Qld only	80 mL

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**IN TASMANIA FOR BLACKBERRY: DO NOT** treat bushes carrying mature or near mature fruit.

**FOR NATIVE VEGETATION:** Use of RELYON TRICLOPYR 600EC HERBICIDE on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

**WITHHOLDING PERIODS:**

**PASTURE: NOT REQUIRED WHEN USED AS DIRECTED.**

**SORGHUM: NOT REQUIRED WHEN USED AS DIRECTED.**

#### GENERAL INSTRUCTIONS

##### COMPATIBILITY

FALLOW SITUATIONS:

RELYON TRICLOPYR 600EC HERBICIDE is compatible with the following products:

- Clopyralid 300 Herbicide
- Fluroxypyr 200 Herbicide
- Glyphosate 450 Herbicide
- Glyphosate-Trimesium 600 Herbicide
- 2,4-D Amine or 2,4-D Ester glyphosate
- Chlorpyrifos 500 EC Insecticide

When mixing with Panzer 450 in fallow, refer to the Panzer 450 label for use rate and adjuvants recommended. DO NOT use Uptake\* Spraying Oil or D-C-Trate Crop Oil.

SORGHUM:

RELYON TRICLOPYR 600EC HERBICIDE is compatible with:

- Fluroxypyr 200 Herbicide
- atrazine (500 g/L flowable product only)

PVC gloves are not recommended for RELYON TRICLOPYR 600EC HERBICIDE, therefore, when tank mixing with products that need to be handled with PVC gloves, workers should wear nitrile gloves.

##### MINIMUM RECROPPING PERIODS

Before using RELYON TRICLOPYR 600EC HERBICIDE in tank mixes with other herbicides, check the plant-back information on all product labels. The time between spraying and planting will be determined by the product with the longest plant-back period.

CRITICAL COMMENTS
<b>DO NOT</b> add crop oils, as severe crop damage may occur. Fusing of sorghum prop roots may be observed and this may be worse under stress (e.g. root disease). Should only be mixed with Fluroxypyr 200 Herbicide and atrazine (500 g/L flowable only) for increased weed spectrum.

Observe the following recropping periods for RELYON TRICLOPYR 600EC HERBICIDE:

wheat, barley, sorghum, maize	7 days
chickpeas, soybeans, sunflowers	7 days
cotton	14 days

##### MIXING

- Half fill the spray unit with water and add the required amount of RELYON TRICLOPYR 600EC HERBICIDE. Add the remaining water with the agitator running. If required, then add crop oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.
- **Basal Bark and Cut Stump Application:** Quarter fill the spray unit or mixing container with diesel and add the required amount of RELYON TRICLOPYR 600EC HERBICIDE. Add the remaining diesel and shake or agitate thoroughly to mix the contents. Periodically shake or agitate to stop product settling out. Only mix sufficient solution for immediate daily use and avoid storing.

##### APPLICATION

###### 1. WOODY WEED SITUATIONS

**Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow to one metre high in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.**

###### A. High Volume Spraying

- Thorough coverage of foliage and stems to the point of runoff is essential, however, avoid excess spraying which is wasteful of chemical.

###### Hand Gun

- Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1500 kPa.
- A spray volume of 3,000 to 4,000 L per infested hectare (30 to 40 L/100 m<sup>2</sup>) should be used on the weed infestation.

###### Knapsack

- Used for smaller infestations, where penetration of the canopy is not essential.
- A spray volume of 3 to 4 L/10 m<sup>2</sup> of infested area should be used.

## B. Aerial Application

- Apply in 100 to 200 L water/ha. Use a calibrated aircraft to apply in half overlap passes. Nozzle configurations should produce droplets of not less than 250 to 350 micron diameter (e.g. D8/45).
- The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets.
- DO NOT spray when wind exceeds 10 km/hr and/or air temperature reaches 30°C.
- Human flagging is not authorised unless protected by engineering controls such as vehicles with cabs.

## C. Controlled Droplet Application (C.D.A.)

- Results similar to high volume spraying can be obtained using Micron Herbi® or similar equipment. Select a nozzle to give a flow rate of 2 mL/sec and sweeping action of approximately 1 m/sec to ensure a droplet density of 20/cm<sup>2</sup>. Use a marking agent as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided with C.D.A. unit.

## D. Low Volume High Concentrate Application Techniques

- Good control will be achieved, similar to high volume application, where bush size enables good coverage of the bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.
- **Gas powered gun:** Apply 50 mL shots to obtain uniform coverage of 4 to 5 m<sup>2</sup> of surface area of bush. This relates to 20 droplets/cm<sup>2</sup> of leaf surface.
- **Sprinkler sprayer:** This technique involves using a micro sprinkler which is connected to a hollow fibre glass rod attached to a pressure knapsack sprayer. Use at low pressures (50 to 200 kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

## E. Basal Bark and Cut Stump Treatment

Mix RELYON TRICLOPYR 600EC HERBICIDE in diesel. The use of diesel as a herbicide carrier may affect the rubber seals in some sprayers. To avoid this, use sprayers which use Viton® seals and fittings. When using RELYON TRICLOPYR 600EC HERBICIDE with diesel, nitrile gloves should be worn instead of rubber gloves.

### Basal Bark Method

- DO NOT apply to wet stems as this can repel the diesel mixture.
- Apply only with hand-directed equipment such as a pressure sprayer or a paint brush.
- Spray equipment should be used at low pressures, up to 200 kPa, to avoid excessive splashing or drift.
- Species with old, rough bark require more thorough wetting than smooth barked species.
- Liberally spray or paint the bark around the stem from ground level up to 30 cm high, wetting thoroughly to the point of runoff (unless otherwise stated).

### Cut Stump Method

- Stems should be cut less than 15 cm above the ground.
- Immediately apply RELYON TRICLOPYR 600EC HERBICIDE/diesel mixture liberally to the freshly cut stump by spray or painting the cut surface and sides of the stem.

## 2. CROPPING/FALLOW SITUATIONS

### A. Boom Application

- Application of RELYON TRICLOPYR 600EC HERBICIDE in a minimum spray volume of 50 L/ha is recommended. Flat fan nozzles are recommended, using pressures in the range of 200-300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

## B. Aerial Application

- **DO NOT** allow RELYON TRICLOPYR 600EC HERBICIDE to physically drift onto desirable plants.
- Aircraft may be used to apply RELYON TRICLOPYR 600EC HERBICIDE in fallow situations, when ground application equipment cannot be used due to prolonged wet conditions.
- A minimum spray volume of 35 L/ha should be used with flat fan and CP nozzle configurations.
- **DO NOT** apply RELYON TRICLOPYR 600EC HERBICIDE by aircraft when wind exceeds 10 km/hr and/or air temperature reaches 30°C.
- Droplets with an average size (or Volume Mean Diameter) of 250 to 350 micron diameter are recommended.
- Human flagging is not authorised unless protected by engineering controls such as vehicles with cabs.

## CLEANING SPRAY EQUIPMENT

### WATER-BASED CLEANING

#### Rinsing

- After using RELYON TRICLOPYR 600EC HERBICIDE, empty the spray unit completely and drain the whole system. Thoroughly wash inside the spray unit using a pressure hose. Drain the spray unit and clean any filters in the tank, pump, lines, hoses and nozzles.
- After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

#### Decontamination

- Before spraying cotton and other sensitive crops, with equipment that has been used to apply RELYON TRICLOPYR 600EC HERBICIDE, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section.
- Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. liquid SURF®, OMO®, DRIVE® at 500 mL/100 L of water or the powder equivalent at 500 g/100 L of water) and circulate throughout the system for at least 15 minutes.
- Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

**Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and water courses.**

### DIESEL-BASED CLEANING

#### Rinsing

- After using RELYON TRICLOPYR 600EC HERBICIDE, empty the spray unit completely and drain the whole system. Thoroughly wash inside the spray unit using a pressure hose. Drain the spray unit and clean any filters in the tank, pump, lines, hoses and nozzles.

On completion of spraying, use a degreaser such as Caltex Kwik-D-Grease® to remove traces of diesel from the sprayer. Rinse tank and spray through the nozzles with water to remove degreaser.

#### Decontamination

- After the above, quarter fill the tank with clean water and add an alkali detergent (e.g. (e.g. liquid SURF®, OMO®, OMOMATIC®, DRIVE® at 50 mL/10 L of water or the powder equivalent at 50 g/10 L of water). Shake or operate spray to circulate the washing solution throughout the sprayer and spray the solution through the nozzle. Rinse well with clean water to remove detergent.

- To clean brushes and container, spray liberally with degreaser. Hose off thoroughly with clean water and repeat using detergents (see above).
- **DO NOT** use this equipment for any other purpose.

#### RESISTANT WEEDS WARNING

### GROUP I HERBICIDE

RELYON TRICLOPYR 600EC HERBICIDE is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group I herbicide.

Some naturally-occurring weed biotypes resistant to the product and Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from the failure of RELYON TRICLOPYR 600EC HERBICIDE to control resistant weeds.

#### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

- Crops susceptible to RELYON TRICLOPYR 600EC HERBICIDE include, but are not limited to: peas, lupins, lucerne, navy beans, soybeans and other legumes; cotton, fruit, hops, ornamentals, shade trees and Pinus spp, potatoes, safflower, sugarbeet, sunflower, tobacco, tomatoes, vegetables and vines.
- RELYON TRICLOPYR 600EC HERBICIDE is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment.
- DO NOT allow physical spray drift onto waterways, native vegetation and susceptible crops. When using RELYON TRICLOPYR 600EC HERBICIDE and Glyphix 450 by aerial application in fallow situations, observance of a buffer zone of 150 metres to protect native tree species is required.
- DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

#### PROTECTION OF LIVESTOCK

- Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

- This product is highly toxic to fish and other aquatic organisms. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

#### STORAGE AND DISPOSAL

Store in closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store near food, feedstuffs, fertilisers or seed.

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**Refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

#### SAFETY DIRECTIONS

Poisonous if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length rubber gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

#### SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

#### CONDITIONS OF SALE

The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

# RELYON®

Nutrien Ag Solutions Limited  
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW 2113.  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92  
relyonaustralia.com.au

®RELYON is a Registered Trademark of Ruralco Holdings Limited