

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TRICLOPYR/PIC

HERBICIDE

ACTIVE CONSTITUENTS:

300g/L TRICLOPYR

present as the BUTOXYETHYL ESTER

100g/L PICLORAM

present as HEXYLOXYPROPYLAMINE SALT

GROUP I HERBICIDE

For control of a range of environmental and noxious woody and herbaceous weeds as specified in the Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL
SHAKE WELL BEFORE USE

RELYON

APVMA APPROVAL NO:
89614/125425

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.

DO NOT spray if rain is likely within one hour or if foliage is wet from rain or dew.

DO NOT burn off, cut or clear blackberry or other woody weeds for at least six months after spraying.

DO NOT apply by aerial application in wind in excess of 15km/hr and air temperatures above 35°C.

In areas prone to flooding treatment should commence after any annual flooding as such areas flooded within 9 months following application may have reduced results.

1 WOODY WEED SITUATIONS

Table A: High Volume Spraying

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ 100L WATER
African boxthorn	Less than 2m tall	All States	500mL
<i>Angophora</i> spp.	1 to 3m tall		350mL
Australian blackthorn	Less than 2m tall		500mL
<i>Banksia</i> spp.	1 to 3m tall		350mL
Biddy bush (Chinese shrub) (Sifton bush)	Autumn when actively growing	ACT and NSW only	500mL
Blackberry in association with: Docks Ragwort Smartweed Thistles	Late spring to autumn	All States	350mL OR 500mL
Blue heliotrope	Flowering	NSW and Qld only	500mL
Brooms: Cape English Flax leaf Montpellier	Spring to mid-summer prior to pod formation	All States	250mL
	Autumn to winter		350mL
Camphor laurel	Less than 2m tall	All States	
	Above 2m tall		500mL
<i>Casuarina</i> spp.	1 to 3m tall		350mL

CRITICAL COMMENTS
Apply when bushes have good leaf cover, growth and no leaf fall .
Apply from late spring to early autumn.
Add a 100% concentrate non-ionic surfactant at 125mL/100L of water for best results.
Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill blackberry. Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Relyon Tricopyr/Pic Herbicide.
Apply in a minimum spray volume of 1250L/ha.
Apply as thorough foliage spray.

Table A: High Volume Spraying

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ 100L WATER
Chinese apple	Less than 2m tall	Qld and WA only	350mL
Cockspur thorn	Spring to autumn	ACT, NSW and Qld only	
Common sensitive plant	Any time when actively growing	NT, Qld and WA only	200mL
Crofton weed	Spring to autumn	ACT, NSW and Qld only	350mL
Eastern cotton bush (<i>Maireana microphylla</i>)	Spring to autumn	NSW and Qld only	500mL
<i>Eucalyptus</i> spp.	Seedling and regrowth from small lignotubers, 1 to 3m tall	All States	350mL OR 500mL
Galenia	Fresh growth in spring to summer	NSW only	500mL
Giant bramble	Spring to autumn	NT, Qld and WA only	
Gorse	1 to 1.5m tall	All States	250mL
	Over 1.5m tall or autumn treatment		350mL
	Winter treatment		500mL
Groundsel bush (<i>Baccharis halimifolia</i>)	1 to 1.5m tall in spring and summer	All States	250mL
	Over 1.5m tall or autumn treatment		350mL

CRITICAL COMMENTS
Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Apply as a thorough foliage spray.
To avoid leaves closing during application, spray plants while moving forward. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Apply as a thorough foliage spray.
Add Uptake Spraying Oil at 500mL/100L of water. Some bushes may require a follow-up spray to control regrowth.
Apply the high rate where difficult to control species of <i>Eucalyptus</i> regrowth is present. Addition of an adjuvant may improve results – contact Relyon for details.
Use 2000L of spray mixture/ha.
Penetration of thick clumps may be difficult and respraying may be necessary. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Spring and Summer treatment only. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Brownout may not be complete until summer. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Apply as a thorough foliar spray.

Table A: High Volume Spraying

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ 100L WATER
Green cestrum	Late spring to early autumn	ACT, NSW and Qld only	500mL
Hawthorn	Less than 2m tall	All States	350mL
Horehound	Pre-flowering	NSW and Qld only	
Japanese sunflower			
Lantana (<i>Lantana camera</i>) (<i>Lantana montevidensis</i>)	Up to 1m tall in summer to autumn	All States	500 or 750mL
	1 to 2m tall in summer to autumn		
Lion Tail (<i>Leonatis nepetifolia</i>)	Pre-flowering	Qld only	200mL
Limebush	Any time of year with good leaf cover and soil moisture	NT and Qld only	350mL
Manuka	At flowering	Vic only	500mL
Mesquite (<i>Prosopis</i> spp.) <i>Prosopis velutina</i>	Seedling, full leaf and flowering before podding	NSW, NT, Qld and WA only	350mL
		Qld only	670mL
Mistflower	Spring to autumn	ACT, NSW and Qld only	350mL
Mother-of-millions	Flowering	NSW and Qld only	500mL
Paddy's lucerne	Active growth	NSW only	
Parkinsonia	Under 2m tall	NT, Qld and WA only	350mL
Prickly pear (common) Smooth tree pear	Active phyllode growth	All States	500mL

CRITICAL COMMENTS
One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1 metre high.
Apply from late spring to early autumn.
Apply as a thorough foliar spray.
Add one of the following adjuvants when using 350mL rate: Uptake Spraying Oil @ 0.5% v/v. Pulse®
Thoroughly wet foliage, stems and soil around the base of the plants. Use higher rate on known harder to kill varieties.
Apply as a thorough foliar spray. Add a 100% concentrate non-ionic surfactant at 100mL/100L for best results.
Penetration of thick clumps may be difficult and respraying may be required. Addition of an adjuvant may improve results – contact Relyon for details.
For optimum results, add Pulse Penetrant at 200mL/100L of spray. Thoroughly wet foliage, stems and soil around the base of the plants. DO NOT spray plants bearing pods. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Apply as a thorough foliar spray.
Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur.
Add Uptake Spray Oil at 500mL/100L water. Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage.
Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.

Table A: High Volume Spraying

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ 100L WATER
Rubber vine (Not infected with rust)	Up to 1.5m tall at flowering	NT, Qld and WA only	350mL
	Dense stands greater than 1.5m tall at flowering		500mL
Siam weed	Active growth	Qld and WA only	350mL
Sicklepod	Up to flowering	NT, Qld and WA only	200mL
St John's wort	From flowering to early seed set	All States	500mL
Sweet briar	Up to 1.5m tall	All States	350mL
Tobacco weed	Actively growing plants	NT, Qld and WA only	300mL
Wattle (<i>Acacia</i> spp.) (except corkwood wattle)	1 to 3m tall	All States	350mL
Wild rosemary (<i>Cassinia laevis</i>)	Active growth, 0.5 to 1m tall	Qld only	350 to 500mL
Wild tobacco tree	Spring to autumn up to 2m tall	ACT, NSW and Qld only	350mL

CRITICAL COMMENTS
Spray all leaves and stems just to the point of runoff and thoroughly spray the base of the plant. With larger, more dense stands regrowth may occur. Subsequent control of any regrowth should be done by basal bark spraying.
Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
DO NOT apply to podding plants. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Late spring to early summer.
Add a 100% concentrate non-ionic surfactant (eg BS 1000) at 100mL/100L of water for best results. Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.
Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Use lower rate on seedlings 0.5m tall. Apply as a thorough foliar spray.

Table B: Aerial Application

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ ha
Blackberry	Summer to autumn	NSW, Qld, SA and Vic and WA only	10L
Grorse		Tas only	
Cockspur thorn Crofton weed Lantana Mistflower	Late autumn	NSW, NT and Qld only (helicopter only)	1.5L plus 7.5L 2,4-D amine (500g/L formulation)
Lantana			10L
Rubber vine (Not infected with rust)	When flowering	NT and Qld only (helicopter only)	3L to 5L
St John's wort	Flowering to early seed set (Nov-Jan)	NSW only	4L
Parkinsonia	Seedlings, 1-2m tall or 12-24 months old	Qld and NT only (helicopter only)	3L

Table C: Controlled Droplet Application (CDA)

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ ha
Blackberry In association with: Docks Ragwort St John's wort Thistles	Summer to autumn	All States except NT	Apply undiluted

CRITICAL COMMENTS
Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Relyon Triclopyr/Pic Herbicide. WARNING: <i>Eucalyptus</i> species up to 4m may be killed if sprayed during this treatment. Mature trees which are 15 to 20m tall may be partially defoliated but are likely to recover.
Helicopter application only.
Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150L/ha. Follow up respraying will be required.
Helicopter application only.
Use rates will depend upon the density and height of the rubber vine stand. The higher rates should be used on dense stands, however, complete coverage and penetration may be difficult. Follow up respraying will be required. Any regrowth should be sprayed with a suitable basal bark herbicide.
Helicopter application only. Follow up spraying will be required in the following season.
Add Uptake Spraying Oil at 1L/ha.

CRITICAL COMMENTS
One application may give satisfactory control but subsequent regrowth and seedlings should be resprayed after hardening off. Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Relyon Triclopyr/Pic Herbicide.

Table D: Low Volume High Concentrate Application Techniques (Gas Powered Gun, Sprinkler Sprayer)

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ 10L water
Blackberry	Late spring to autumn	ACT, NSW, Qld, SA, Tas and WA only	335mL
Camphor laurel Cockspur thorn Crofton weed	Less than 1.5m high	ACT, NSW and Qld only	500mL
<i>Eucalyptus</i> species	Seedlings up to 2m tall	All States	335mL
Mistflower	Less than 1.5m high	ACT, NSW and Qld only	500mL
Sweet briar	1.5m tall, full leaf to ripe fruit	NSW only	
St John's wort	During flowering to early seed set	NSW, Tas and Vic only	
Wild tobacco tree	Less than 1.5m high	ACT, NSW and Qld only	

Table E: Boom Application

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ ha
Galenia	Fresh growth during spring to summer	NSW only	5L
Sicklepod	Up to flowering	NT and Qld only	3L
St John's wort	Flowering to early seed set (Nov-Jan)	NSW only	2 to 4L

CRITICAL COMMENTS
Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.
Gas Powered Gun only: Apply to actively growing bushes not more than 1.5m tall that have not more than 5 stems from the crown.
Gas Powered Gun only: One application should provide control. Minor regrowth and seedlings may be retreated the following summer.
Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.

CRITICAL COMMENTS
Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.
DO NOT apply to podding plants. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water.
Use the higher rate on dense infestations and when longer residual control is required.
Follow up respraying will be required in the following season.

2 FALLOW SITUATIONS

Table A: Boom Application

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ ha	
Blackberry nightshade – Suppression only	10 to 25cm tall, prior to flowering	NSW and Qld only	200 to 400mL + 1.2L Glyphosate 450 g/L Herbicide + Adjuvant	
Camel melon Prickly paddy melon Cucumber melon (<i>Cucumis melo</i>)	From 2 leaf to 50cm diameter			
Common sowthistle	From 8 leaf to flowering			
Cow vine	From 2 to 5 leaf up to 15cm diameter, prior to flowering			
Lucerne (established)	Active growth, 15 to 25cm high, during spring			300 to 500mL + 1.2L Glyphosate 450 g/L Herbicide + Adjuvant
<i>Polymeria pusilla</i>	2 to 12 leaf up to 20cm diameter, prior to flowering			200 to 400mL + 1.2L Glyphosate 450 g/L Herbicide + Adjuvant

CRITICAL COMMENTS
<p>FOR USE BY GROUND EQUIPMENT ONLY.</p> <p>Plants must be actively growing.</p> <p>Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5cm diameter for <i>Polymeria pusilla</i>).</p> <p>Refer to Glyphosate 450 g/L herbicide label for use of adjuvant.</p> <p>DO NOT plant susceptible crops for up to nine months after application, as specified in General Instructions - Minimum Recropping Periods - Black Cracking Clay Soils, NNSW & Qld.</p> <p>Dry conditions after application will increase the recropping interval.</p>

Table B: Blanket Wiper Application

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

FALLOW			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE
Bitterbark (<i>Alstonia constricta</i>)	From summer to end of autumn	Qld	1:4 (1 part Relyon Triclopyr/Pic Herbicide to 4 parts water)
			2% solution for spot spraying (eg 100mL Relyon Triclopyr/Pic Herbicide in 5L water)

CRITICAL COMMENTS
<p>For use with blanket wipers only.</p> <p>For best results apply in autumn to tall (> 60cm) plants using two opposite directional passes (up and back). Follow up “missed” plants with a spot spray application. These will be obvious after 6 weeks.</p> <p>Blanket wiper applications can be made in summer when plants are smaller but follow up spraying may be necessary.</p> <p>Do not disturb (cultivate) the treated patches for at least 3 months. Best long term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months).</p> <p>Spot spraying “missed” plants: Thoroughly wet all stems and leaves without producing any solution run-off. Avoid any spray reaching the soil surface.</p>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

IN TASMANIA FOR BLACKBERRY

DO NOT treat bushes carrying mature or near mature fruit.

FOR NATIVE VEGETATION

Use of Relyon Triclopyr/Pic Herbicide on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

WITHHOLDING PERIOD

NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

MINIMUM RECROPPING PERIODS – Black Cracking Clay Soils, NNSW & Qld.

Table A: Boom Application

Plant-back periods for crops following the application of Relyon Triclopyr/Pic Herbicide for rates up to 600mL/ha.				
RATE mL/HA	200	300	400	600
CROP	Months			
Wheat	2	2	4	4
Barley	2	2	4	4
Canola	2	4	4	4
Faba bean	4	4	6	6
Chickpea	4	6	6	6
Lucerne	6	9	9	9

These plant-back periods are based on a normal rainfall pattern. During drought conditions (or when rainfall is less than 100mm for a period of 4 months or greater) the plant-back period may be significantly longer.

Table B: Blanket Wiper Application

Plant-back periods for crops following blanket wiper application.	
CROP	Months
Broadleaf Crops	18
Lucerne	6
This will allow any potential soil residues to dissipate, if any, and allow effective control.	

Note: Before using Relyon Triclopyr/Pic Herbicide in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product, ie. the product with the longest plant-back period, will determine the time between spraying and planting.

COMPATIBILITY

Follow any regional restrictions, and all directions and restrictions on the label, of any chemical mixed with Relyon Triclopyr/Pic Herbicide (eg. 2,4-D amine).

Relyon Triclopyr/Pic Herbicide is compatible with the following herbicides:
2,4-D Amine, DP® 600, Metsulfuron methyl, Glyphosate 450 g/L, Fluroxypyr 200 g/L, Glyphosate 490 g/L

Relyon Triclopyr/Pic Herbicide is compatible with the following adjuvants, as per Directions for Use:
Uptake, Pulse, non-ionic surfactant (1000g/L)

MIXING

Mix only with water.

Half fill the spray unit with water, and add the required amount of Relyon Triclopyr/Pic Herbicide. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

APPLICATION

1 WOOD WEED SITUATIONS

Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow to approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.

A High Volume Spraying

- Thorough coverage of foliage to the point of run-off is essential, however, avoid excess spraying which is wasteful of chemical.

Hand Gun

- Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1500kPa (400 to 500kPa for St John's wort).
- A spray volume of 3000 to 4000L per infested hectare of 1 to 2 metre high blackberry (30 to 40L/100m²) should be used.
- Use 2000L of spray mixture/ha of Galenia infestation (ie. 20L/100m² infested area).

Knapsack

- Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to hand gun.
- A spray volume of 3 to 4L/10m² infested area should be used.
- A spray volume of 2L/10m² should be used for an area infested with Galenia.

B Aerial Application

- Apply in 200L of water/ha using an aircraft to apply 100L per pass on a double overlap pattern using nozzle configurations to produce droplets of 250 to 350 micron diameter.
- The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets.
- **DO NOT** spray when wind exceeds 15km/hr and/or air temperature reaches 35°C.

C Controlled Droplet Application (CDA)

- Results similar to high volume spraying can be obtained used Micron Herbi® or similar equipment. Select a nozzle to give a flow rate of 2mL/sec and sweeping action of approximately 1m/sec to ensure a droplet density of 20/cm². Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided by CDA unit.

D Low Volume High Concentrate Application Techniques

- Good control will be achieved, similar to high volume application, where bush size enables good coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.
- **Gas Powered Gun:** Apply 50mL shots to obtain uniform coverage of 4 to 5m² of surface area of bush. This relates to 20 droplets/cm² of leaf surface.
- **Sprinkler Sprayer:** This technique involves using a micro sprinkler that is connected to a hollow fibreglass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50 to 200kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

E Boom Application

- Application in a minimum spray volume of 200L/ha for Galenia and St John's wort and 600L of water/ha for Sicklepod. Flat fan nozzles are recommended, using pressure in the range of 200-300kPa. Boom height must be set to ensure double overlap of nozzle patterns.

2 FALLOW SITUATIONS

A Boom Application

- Application of Relyon Triclopyr/Pic Herbicide in a minimum spray volume of 50L/ha is recommended. Flat fan nozzles are recommended, using pressure in the range of 200-300kPa. Boom height must be set to ensure double overlap of nozzle patterns.

B Blanket Wiper Application

- Blanket needs to be made from durable and wettable material with a rigid backing.
- Blanket should be rigidly mounted behind motorised vehicle (eg. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (rewetting).
- Ideally, a scraper bar should be mounted in front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping the stems. This scraper may be mounted at the front of the vehicle.
- Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10-15kph are ideal for blanket wiping application.

CLEANING SPRAY EQUIPMENT

Rinsing

- After using Relyon Triclopyr/Pic Herbicide, empty spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles.

- After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

Decontamination

- Before spraying cotton and other sensitive crops with equipment that has been used to apply Relyon Triclopyr/Pic Herbicide, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section.
- Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (eg. liquid SURF®, OMO®, OMO Matic®, DRIVE® at 500mL/100L of water or the powder equivalent at 500g/100L of water) and circulate throughout the system for at least 15 minutes.
- Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

RESISTANT WEEDS WARNING

GROUP I HERBICIDE

Relyon Triclopyr/Pic Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group I herbicide.

Some naturally occurring weed biotypes resistant to the product and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Nutrien Ag Solutions Limited representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Crops susceptible to Relyon Triclopyr/Pic Herbicide include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and *Pinus* spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines.

Relyon Triclopyr/Pic Herbicide is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment.

Picloram, one of the active constituents in this product, can remain in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation.

DO NOT allow physical spray drift onto waterways, native vegetation and susceptible crops.

DO NOT apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants.

DO NOT apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants.

DO NOT move soil, which may have been treated to areas where desirable plants are to be grown.

PROTECTION OF LIVESTOCK

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.

Many plants remain poisonous after death, and stock should not be allowed access it, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or watercourses with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** store near feedstuffs, fertilisers or seed.

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SMALL SPILL MANAGEMENT

Wear protective equipment (See SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb as above, the wash liquid for disposal.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin.

When preparing the spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

RELYON®

Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW 2113.
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92
relyonaustralia.com.au

®RELYON is a Registered Trademark of Ruralco Holdings Limited